



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

As the weather conditions for many of the fourteen exposures were such as to vitiate the results, six of the plates which are extremely good in quality were on this account measured again, giving:

$$\pi_{\text{rel.}} = -0''.007 \pm 0''.006.$$

The negative parallax derived in both cases indicates that the real parallax must be small, probably less than $+0''.005$. This result is quite remarkable in relation to the motion (in one case reaching 20 to 30 seconds of arc annually) found by Hubble for some parts of the nebula¹. Assuming as high a velocity as 1000 km/sec., this motion would still require a parallax of about $0''.100$, which is quite inconsistent with our present results. It seems necessary, therefore, to assume that the motions are a question of illumination, in which case we have to deal with the velocity of light. As we do not, however, know the orientation of the nebula, we can only conclude that the parallax must be small.

A. VAN MAANEN.

THE PARALLAXES OF TWO LONG-PERIOD VARIABLES

Eighteen exposures each of *T Cassiopeiae* and *R Trianguli* give as their parallaxes:

$$\text{T Cass.} \quad \pi_{\text{rel.}} = +0''.027 \pm 0''.003$$

$$\text{R Triang.} \quad \pi_{\text{rel.}} = +0''.005 \pm 0''.005$$

For only one other star of this type, *o Ceti*, is a parallax available: Russell gives $\pi_{\text{rel.}} +0''.136$; Kostinsky, $+0''.013$; Mitchell, $+0''.061$. Applying the systematic corrections given in *Mount Wilson Contribution* No. 189, the absolute parallaxes of the three objects are $+0''.027$, $+0''.005$, and $+0''.047$. Using these values, we obtain the absolute magnitudes, *M*, at maximum and minimum, given in the following table:

	m		M	
	Max.	Min.	Max.	Min.
T Cass.....	6.7	12.5	+3.9	+9.7
R Triang.....	6.5	12.0	0.0	+5.5
o Ceti.....	1.7	9.6	+0.1	+8.0

A. VAN MAANEN.